

Superfunctions and sqrt of Factorial

D. Kouznetsov^{1a}, H. Trappmann^b

¹ *Institute for Laser Science, University of Electro-Communications 1-5-1 Chofugaoka,
Chofushi, Tokyo, 182-8585, Japan*

² *Kameruner Str. 9, 13351 Berlin, Germany*

E-mail: ^a dima@ils.uec.ac.jp , ^b henryk@pool.math.tu-berlin.de

The English version compiled 10 сентября 2009 г.

The holomorphic function h is constructed such that $h h z = z!$; this function is interpreted as square root of Factorial.

Keywords: sqrt of Factorial, superfunction, SuperFactorial, inverse problem

PACS: 02.30.Ks,02.30.Zz,02.30.Gp,02.30.Sa,

Introduction

This work was motivated by one exercise from the past-century course of Quantum Mechanics of the Moscow State University. It was suggested to give sense to the operator $\sqrt{!}$ [1]. That time, a satisfactory solution was not found; the opinion was, that such an operation has no meaning [2].

In Quantum Mechanics, the repeated (iterated) application of an operation (usually, some “observable”) to some argument (which may have sense of vector of state) is interpreted as “power” of the operation; in particular, in such a way the square of coordinate or the square of momentum are treated. For this analogy, the notation without parenthesis is used. In these notations, $\sin \alpha$ means $\sin(\alpha)$, $\ln \sin x$ means $\ln(\sin(x))$ and so on; such notations are used also in textbooks on elementary algebra. To avoid confusions at the iterations, we use also the prefix notation $\text{Factorial } z = \text{Factorial}(z)$ instead of $z!$.

We assume, that factorial is known meromorphic function, just Gamma function [3] with displaced argument. In this way factorial is interpreted in programming languages Mathematica and Maple. The factorial of the real argument is shown in figure 0.

In this work, the square root of the Factorial is interpreted as a holomorphic function h such that its second iteration is Factorial, i.e. $h h z = z!$. For real values of argument the graphic of function $\sqrt{!}$ is shown in figure 1. Below we describe, how to evaluate it not only for real, but also for complex values of the argument, using superfunctions [4–6].

1. Superfunctions

The evaluation of fractional power of a function, id est, the fractional iteration, (for example,

$\sqrt{\exp}$, see [7–9], or $\sqrt{!}$), can be based on the concept of superfunction [4–6]. For a given function H , which is referred to as the “Transfer Function” below, a superfunction F is a holomorphic solution of the Abel equation

$$F(z+1) = H(F(z)) \quad . \quad (1)$$

Such an equation is pretty old [10–12], although in 1827, N.H.Abel wrote it in a different form, for the inverse function of F . The Abel equation comes from the phenomenological consideration of the transformation of a signal F in a single-dimensional homogenous nonlinear system, characterized with the Transfer Function H . Equation (1) may have also other applications, discussed below in the special section. In some sense, the equation (1) is equivalent of the Schröder’s equation [13–17]; at the exponential transformation of the argument, the inverse Schröder’s function becomes a superfunction, but not every supefunction can be simply expressed through some inverse Schröeder function. For this reason, here we deal with superfunctions and not with the Schröder functions.

A superfunction F determines the fractional iteration H^c of the Transfer Function H :

$$H^c(z) = F(c + F^{-1}(z)) \quad . \quad (2)$$

The resulting function H^c can be considered as fractional power, of function H , because it satisfies the expected relations

$$H^1 = H \quad , \quad H^{c+d}(z) = H^c H^d z = H^c(H^d(z)) \quad ,$$

id est, for two numbers c and d , the identity $H^c H^d = H^{c+d}$ holds, as if H would be not a function but a number. In particular with $c=1/2$, the half-iteration $h(z) = \sqrt{H}(z) = H^{c}(z)$ is considered to be the square root of function H , because $h h z = h(h(z)) = H z = H(z)$. In this sense $h^2 = H$ and $h = \sqrt{H}$.

Some superfunctions (see Table 1) are well known; they are used without to identify them as superfunctions. Several elementary superfunctions (in particular, trigonometric) were listed also at websites [5, 6, 18].

Superfunctions of the exponential (see row 4 of the Table 1) yet are not so widely known, although Helmuth Kneser had reported the half-iteration of the exponential, id est, $\sqrt{\exp}$, in year 1950 [7]. Here, tetrational tet is the superfunction of exp, characterized in that $\text{tet}(0) = 1$ and holomorphic on the range $\mathbb{C} \setminus \{x \in \mathbb{R} : x \leq 0\}$; id est, holomorphic solution of equation

$$\text{tet}_b(z+1) = \exp_b(\text{tet}_b(z)). \quad (3)$$

At integer values of z , tetration $\text{tet}_b(z)$ is result iterational applicaiton of exponential to unity:

$$\text{tet}_b z = \underbrace{\exp_b \left(\exp_b \left(\dots \exp_b(1) \dots \right) \right)}_{z \text{ repetitions of exponential}} \quad (4)$$

TABLE 1. Examples of superfunctions

#	$H(z)$	$F(z)$	$F^{-1}(z)$	comment
1	$z+1$	$b+z$	$z-b$	$b \in \mathbb{C}$
2	$b+z$	$bz+c$	$(z-c)/b$	$b \neq 0$
3	$bz+c$	$b^z + \frac{c}{1-b}$	$\log_b(z - \frac{c}{1-b})$	
4	b^z	$\text{tet}_b(z)$	$\text{tet}_b^{-1}(z)$	(3),(4), [4, 9, 19–22]
5	z^b	$\exp(b^z)$	$\log_b(\ln(z))$	$b > 0$
6	$\ln(b+e^z)$	$\ln(bz)$	e^z/b	$b \neq 0$
7	$(a^b+z^b)^{1/b}$	$az^{1/b}$	$(z/a)^b$	$a > 0, b \neq 0$
8	$2z^2-1$	$\cos(\pi 2^z)$	$\log_2(\arccos(z)/\pi)$	
9	$2z^2-1$	$\cosh(\pi 2^z)$	$\log_2(\text{arccosh}(z)/\pi)$	compare #8
10	$2z/(1-z^2)$	$\tan(\pi 2^z)$	$\log_2(\arctan(z)/\pi)$	
11	$2z/(1+z^2)$	$\tanh(\pi 2^z)$	$\log_2\left(\frac{2}{\pi} \ln\left(\frac{z+1}{z-1}\right)\right)$	
12	Factorial(z)	SuperFactorial(z)	ArcSuperFactorial(z)	(6),(8)
	$P(H(Q(z)))$	$P(F(z))$	$F^{-1}(Qz)$	$P(Q(z))=z$

Evaluation of tet_b at $b > \exp(1/e)$ and in particular, for $b = e$ and $b = 2$ is described in [9, 19, 22]. For $b = e$ the fast approximation is available at [20, 21]. The evaluation at $1 < b < \exp(1/e)$ and, in particular, for $b = \sqrt{2}$, is considered in [4].

New superfunctions can be obtained by transformation of the already established superfunctions. If some F is superfunction of some Transfer Function H , then another superfunction \mathcal{F} can be defined as $\mathcal{F}(z) = F(z + \delta(z))$, where δ is some holomorphic 1-periodic function. In particular, the superfunction in row 9 can be obtained from that in row 8 with $\delta(z) = \pi \cdot i \cdot \ln(2)/2 = \text{constant}$. For more complicated function δ , the resulting function \mathcal{F} usually has either a reduced range of holomorphizm, or some fast growth in the direction of the imaginary axis.

In addition, for the pair of the mutually-inverse functions P and Q , the new pair (Transfer Function, superfunction) can be obtained by the transform, indicated in the last row of the Table 1, from the previous rows. Therefore, the table 1 can be much larger.

Table 1 can be extended also considering any pair F and F^{-1} of biholomorphic functions, declaring F as superfunction, and constructing the corresponding Transfer Function with (2) at $c = 1$. In this work the inverse problem is considered, id est, the building of the superfunction for the given Transfer Function, namely, Factorial. This allows to evaluate the holomorphic function $\sqrt{!}$, giving sense to the logo of the Physics department of the Moscow State University.

2. Factorial, ArcFactorial and the fixed points

For the evaluation of superFactorial and its inverse, efficient implementations of Factorial and ArcFactorial are required. Complex maps of Factorial and ArcFactorial are shown in figure 2. In the right hand side of the left picture, the density of levels for the Factorial is so high that they would overlap; this area left empty. We use the original numerical C++ implementation [23] of the Factorial and ArcFactorial.

At the analysis of a superfunction for some Transfer Function H , the key question is about the fixed points [16, 17]. In the case $H = \text{Factorial}$, the fixed points are solutions of equation $\text{Factorial}(z) = z$. The real fixed points correspond to abscissas of the intersecions of the graphics $y = \text{Factorial } x$ and $y = x$. These graphics are shown in figure 0.

The $\text{Factorial}(z)$ has two obvious fixed points $z = 1$ and $z = 2$. However, there is also a countable set of negative fixed points. Two of them are seen in the left bottom corner of figure 0.

Each of the fixed points can be used to build-up the superfunction and corresponding non-integer iterations of the transfer function [4]; there exist many functions that can be considered as a square root of the Factorial. In this work, the superfunction built up at the fixed point $z = 2$, and the corresponding $\sqrt{!}$ is considered. Such a choice is determined by the intent to build-up a function that is holomorphic while the argument is larger than 2, and grow up slower than the factorial, which corresponds to the intuitive expectation about such a function. Namely such a factorial could be used for description of processes that grow up faster than any polynomial but slower than any exponential. Namely this $\sqrt{!}$ is plotted in the left hand side of figure 1 with a solid curve.

3. Evaluation of SuperFactorial

Consider the SuperFactorial $= F$, that approaches the fixed point 2 at large negative values of the real part of the argument:

$$\text{Factorial}(F(z)) = F(z+1) , \quad \lim_{x \rightarrow -\infty} F(x+iy) = 2 . \quad (5)$$

Following the algorithm [4], we search the solution F in the form

$$F(z) = \Phi(\exp(kz)) , \quad (6)$$

where k is constant; function Φ can be interpreted as the inverse Schröder function. The substitution of transformation (6) into equation (5) gives the Schröder equation [13, 14]

$$\Phi(K\varepsilon) = \text{Factorial}(\Phi(\varepsilon)) , \quad (7)$$

Table 2. Coefficients u and U in the expansions (8) and (9).

n	u_n	U_n
2	0.7987318351724345	-0.7987318351724345
3	0.5778809754764832	0.6980641135593670
4	0.3939788096629718	-0.6339640557572815
5	0.2575339580323327	0.5884152357911399
6	0.1629019581037053	-0.5538887519936520
7	0.1002824191713524	0.5265479025985924
8	0.0603184725913977	-0.5041914604280215
9	0.0355544582258062	0.4854529800293392
10	0.0205859954874424	-0.4694346809094714

where $K = \exp(k)$. We search the solution Φ in the form

$$\Phi(\varepsilon) = 2 + \varepsilon + \sum_{n=2}^N u_n \varepsilon^n + \mathcal{O}(\varepsilon^{N+1}), \quad (8)$$

where u are constant real coefficients, and integer $N > 0$. The substitution of such expansion into equation (7) gives the value

$$k = \ln(K) = \ln(3 + 2 \text{Factorial}'(0)) = \ln(3 - 2\gamma) \approx 0.6127874523307,$$

where $\gamma = -\Gamma'(1) \approx 0.5772156649$ is the Euler constant [3], and the chain of equations for u . From this chain we find

$$u_2 = \frac{\pi^2 + 6\gamma^2 - 18\gamma + 6}{12(3 - 5\gamma + 2\gamma^2)} \approx 0.798731835$$

$$u_3 = \frac{-36 - 39\pi^2 - 738\gamma^2 + 324\gamma + 99\pi^2\gamma - 60\pi^2\gamma^2 - \pi^4 + 24\gamma^5 + 594\gamma^3 - 120\zeta(3)\gamma + 48\zeta(3)\gamma^2 + 12\gamma^3\pi^2 + 72\zeta(3) - 204\gamma^4}{144(-18 + 69\gamma - 104\gamma^2 + 77\gamma^3 - 28\gamma^4 + 4\gamma^5)}$$

where $\zeta(z) = \sum_{n=1}^{\infty} \frac{1}{n^z}$ is the Riemann zeta-function [3]; $\zeta(3) \approx 1.202056903$. Similar (but longer) expressions can be obtained for other coefficients u ; few first coefficients are evaluated in Table 2. In addition, we set $u_0 = 2$ and $u_1 = 1$. The partial sum in (8) at $N = 20$ gives the approximation F_{20} for the superFactorial

$$F_{20}(z) = \sum_{n=0}^{20} u_n \exp(knz),$$

providing of order of 15 correct decimal digits at $\Re(z) \leq -2$. For evaluation of $F(z)$ at $\Re(z) > -2$, the recursive formula is used, $F(z) = \text{Factorial}(F(z-1))$. Such numerical representation is used to plot the map of SuperFactorial in the complex plane in figure 3, as well as the real-real plot in figure 1.

The SuperFactorial F is entire periodic function. The period

$$T = 2\pi i/k \approx 10.2534496811560279265772640691397 i$$

is pure imaginary, so, the pattern in the plot of this function in figure 3 is reproduced at the translations along the imaginary axis. Along the real axis, the SuperFactorial F grows up faster, than any exponential (see Fig.1), and even faster than tetration [9].

In vicinity of the real axis, the plot of the SuperFactorial shows the quasi-periodic, “fractal” structure, similar to that of growing superexponentials [4, 22]. Perhaps, other holomorphic functions that grow up faster than any finite iteration of exponential (id est \exp^c at fixed c) behave in the similar way.

4. ArcSuperFactorial

The inverse function of the SuperFactorial, id est, ArcSuperFactorial= $G=F^{-1}$, is shown in the right hand picture of figure 3. This function can be expressed, inverting expansion (8):

$$G(z) = \frac{1}{k} \log \left(\sum_{n=1}^N U_n (z-2)^n + \mathcal{O}(z-2)^{N-1} \right), \quad (9)$$

where $U_1 = 1$ and $U_2 = -u_2$. The Mathematica routine InverseSeries (see also [3],eq.3.6.25) allows to express coefficients U in terms of coefficients u ; both u and U are evaluated in table 2.

The partial sum in (9) at $N=20$ is used as approximation for G , allowing to evaluate the $G(z)$ with 15 decimal digits at $|z-2| \leq 0.1$. For other values of the argument, the recurrent equation can be used: $G(z) = G(\text{ArcFactorial}(z)) + 1$. The iterations of ArcFactorial converge to value 2; after several such iterations, the G can be evaluated by expression (9). With complex < double > arithmetics, of order of 14 significant figures can be achieved in such a way.

ArcSuperFactorial $G=F^{-1}$ is holomorphic function on the domain $\mathbb{C} \setminus \{x \in \mathbb{R} : x \leq 2\}$. Along the real axis, G grows until infinity, although slowly; slower than any logarithm and even slower than ArcTetration [9]. Similar slow growth of the modulus of ArcSuperFactorial takes place at the moving out from the origin of coordinats in any direction, except the negative direction of the real axis. Number 2 is branchpoint of G ; the cut goes left along the real axis.

5. Square root of Factorial

The choice of the SuperFactorial F and ArcSuperFactorial G determines any power of the factorial; in particular, the fractional power and even the complex power. (The Transfer Function can be iterated complex number of times.) If the number of iterations is equal to one half, the combination (2) of SuperFactorial and ArcSuperFactorial gives the $\sqrt{!}$, id est,

the square root of the Factorial.

$$\sqrt{!}(z) = \text{Factorial}^{1/2}(z) = F(1/2 + G(z)) . \quad (10)$$

This function in the complex plane is shown in the left hand side of the figure 4. For comparison, the right hand picture of figure 4 shows the square root of the exponential $\sqrt{\exp}(z) = \text{tet}(1/2 + \text{tet}^{-1}(z))$, where tetrational tet is superfunction of the exponential [9], id est the holomorphic solution of equation (3). The fast numerical implementation of tetrational tet holomorphic in the domain $\mathbb{C} \setminus \{x \in \mathbb{R} : x \leq -2\}$, and arctetration tet^{-1} is described in [20]; the Mathematica code for the evaluation is available at [21]. The Arctetration tet^{-1} may be called also “superlogarithm” [24], although it is not a superfunction of the logarithm.

The constructed $\sqrt{!}$ is a holomorphic function on the domain $\mathbb{C} \setminus \{x \in \mathbb{R} : x \leq 1\}$. In general, $\sqrt{!}$ behaves in a way, similar to $\sqrt{\exp}$; however, it has only one branch point and only one cut line, while the $\sqrt{\exp}$ has two; also, $\sqrt{!}$ grows up a little bit faster, than the $\sqrt{\exp}$. The behavior of $\sqrt{!}$ for real values of the argument is shown in figure 1. In general, this function corresponds to the intuitive expectations about its behavior. In particular, it grows faster than any polynomial, but slower than any exponential. In the similar way, maps of functions Factorial^c and \exp^c can be plotted for other values c . In particular, at $c=1$, this function is Factorial; at $c=-1$, it is ArcFactorial, shown in figure 2; at $c=0$, H^c becomes identical function. In such a way, equation (2) allows the smooth (holomorphic) transition from the Transfer Function to its inverse function. Such transitions can be defined also for other Transfer Functions.

6. Physical applications

Although a distributed physical system with the Factorial as Transfer Function is not reported, the same method can be applied for other Transfer Functions. In this section, the possible applications of the formalizm of superfunctions are discussed.

In the investigation of the nonlinear response of optical materials, the sample is supposed to be optically thin, in such a way, that the intensity of the light does not change much as it goes through. Then one can consider, for example, the absorption as function of the intensity. However, at small variation of the intensity in the sample, the precision of measurement of the absorption as function of intensity is not good. The reconstruction of the superfunction from the Transfer Function allows to work with relatively thick samples, improving the precision of measurements. In particular, the Transfer Function of the similar sample, which is half thinner, could be interpreted as the square root (id est, half-iteration) of the Transfer Function of the initial sample.

In nonlinear acoustics, it may have sense to characterize the nonlinearities in the attenuation of shock waves in a homogeneous tube. This could find an application in some advanced muffler, using nonlinear acoustic effects to withdraw the energy of the sound waves without to disturb the flux of the gas. Again, the analysis of the nonlinear response, id est, the Transfer Function, may be boosted with the superfunction.

In analysis of condensation, the growth (or vaporation) of a small drop of liquid can be considered, as it diffuses down through a tube with some uniform concentration of vapor. In the first approximation, at fixed concentration of the vapor, the mass of the drop at the output end can be interpreted as the Transfer Function of the input mass. The square root of this Transfer Function will characterize the tube of half length.

In a similar way the mass of a snowball, that rolls down from the hill, can be considered as a function of the path it already have passed. At fixed length of this path (that can be determined by the altitude of the hill) this mass can be considered also as a Transfer Function of the input mass. The mass of the snowball could be measured at the top of the hill and at the bottom, giving the Transfer Function; then, the mass of the snowball as a function of the length it passed is superfunction.

If one needs to build-up an operational element with factorial transfer function, and wants to realize it as a sequential connection of a couple of identical operational elements, then, each of these two elements should have transfer function $\sqrt{!}$ shown in figure 1.

Calculation of a fractal iteration of a function, and in particular, that of exponential or that of the Factorial, may have other (and, perhaps, unexpected) applications; for example, in the description of the physical processes, that grow up faster than any polynomial, but slower than any exponential. The theoretical science should be ready for such applications. In particular, the SuperExponential, SuperFactorial, $\sqrt{\exp}$, $\sqrt{!}$ should be popped up to the level of special functions.

Conclusion

A SuperFactorial is constructed as superfunction of the Factorial, solving the equation (5). An arbitrary power c of a function H can be expressed through its superfunction F by (2). For $H = \text{Factorial}$ and $c = 1/2$, this gives way of evaluation of holomorphic function $\sqrt{!}$ shown in figures 1 and 4. The formalism of superfunctions allows the evaluation of non-integer iterations of various functions and may have applications in physics and technology.

Authors are grateful to R.D.Kouznetsov, M.A.Kallistratova, A.V.Borisov and M.Sadgrove for the important crytics, to S.Okudaira for the indications of the new applications of superfunctions, and to participants of the Tetration Forum [6] and Citizendum [5] for

important discussions.

Список литературы

1. I.M.Saraeva, Yu.M.Romanovsky, A.V.Borisov (composers). Vladimir Dmitrievich Krivchenkov. Physical Department of MSU, 2008, (in Russian), online version, p.81; <http://www.phys.msu.ru/rus/about/structure/admin/OTDEL-IZDAT/HISTORY/>
2. Gordienki V.M., Novik V.K. About the time, the department, the laboratory and myself... 70 anniversary of V. P. Kandidov. Moscow, 2007. (In Russian) http://ofvp.phys.msu.ru/pdf/Kandidov_70.pdf
3. Abramowitz M., Stegun I. Handbook on special functions. NY, 1968.
4. Kouznetsov D., Trappmann H. Portrait of the four regular super-exponentials to base $\sqrt{2}$. Mathematics Of Computation, in press. Preprint ILS UEC, 2009: <http://www.ils.uec.ac.jp/~dima/PAPERS/2009sqrt2.pdf>
5. <http://en.citizendium.org/wiki/Superfunction>
6. Trappmann H. Elementary superfunctions. Tetration Forum. Berlin, 2009. <http://math.eretrandre.org/tetrationforum/showthread.php?tid=275>
7. Kneser, H. Reelev analytische Lösungen der Gleichung $\varphi(\varphi(x)) = e^x$. J. f. reine angew. Math. 1950. **187**. P. 56–67.
8. Trappmann H., Kouznetsov D. Uniqueness of holomorphic Abel functions. Preprint ILS UEC, 2009. <http://www.ils.uec.ac.jp/~dima/PAPERS/2009uniabel.pdf>
9. Kouznetsov D. Analytic solution of $F(z+1)=\exp(F(z))$ in complex z -plane. Mathematics of Computation. 2009. **78**. P. 1647-1670.
10. Belitskii G., Lubish Yu. The real-analytic solutions of the Abel functional equations. Studia Mathematica 1999 . **134**. N 2. P. 135–141.
11. Abel N.H. Correlative of the functional equation. Crelle's Journal 1827 N 2. P. 389.
12. http://en.citizendium.org/wiki/Abel_function
13. Sui Sun Cheng, Wenrong Li. Analytic solutions of functional equations. World Scientific, 2008.
14. Schröder E. Über iterierte Funktionen III. Math. Ann. 1870. **3**. P. 296–322.

15. Kuczma M., Choczewski B., Ger R. Iterative Functional Equations Cambridge, 1990.
16. Écalle J. 1970. Théorie des invariants holomorphes. Publications d'mathématiques d'Orsay, 67-74 09. http://portail.mathdoc.fr/PMO/PDF/E_ECALLE_67_74_09.pdf
17. Milnor J., Dynamics in One Complex Variable. Princeton, 2006.
18. Mueller M. Old projects. <http://www.math.tu-berlin.de/~mueller/projects.html>
19. Kouznetsov D. Ackermann functions of complex argument. Preprint ILS UEC, 2008; <http://www.ils.uec.ac.jp/~dima/PAPERS/2008ackermann.pdf>
20. Kouznetsov D. Superexponential as special function. Vladikavkaz Mathematical Journal, in press. Preprint: <http://www.ils.uec.ac.jp/~dima/PAPERS/2009vladie.pdf>
21. <http://en.citizendium.org/wiki/TetrationDerivativesReal.jpg/code>
22. Kouznetsov D., Trappmann H. Cut lines of the inverse of the Cauchi-tetrational. Preprint ILS UEC, 2009. <http://www.ils.uec.ac.jp/~dima/PAPERS/2009fractae.pdf>
23. <http://en.citizendium.org/wiki/ArcGamma.jpg/code>
24. Robbins A. Solving for the Analytic Piecewise Extension of Tetration and the Super-logarithm. Preprint of "Virtual Composer", 2000. http://misc.virtualcomposer2000.com/TetrationSuperlog_Robbins.pdf




Рис. 0: Factorial of the real argument and graphical solution of equation $x! = x$.





FIG. 1: Left: $y = \text{Factorial}(x)$, thick dashed curve; $y = \sqrt{x!} = \text{Factorial}^{1/2}(x)$, solid curve; $y = \sqrt{x!} = \text{Factorial}(x)^{1/2}$, thin dashed, as functions of real x ; Right: $y = F(x)$, solid, compared to $y = 2 + \exp(kx)$, dashed




Рис. 2: $f = \text{Factorial}(z)$, left, and $f = \text{ArcFactorial}(z)$, right, in the complex z -plane; Levels $\Re(f) = p = \text{constant}$ and Levels $\Im(f) = q = \text{constant}$ are shown with thick lines for $p = -4, -3, -2, -1, 0, 1, 2, 3, 4$ and for $q = -4, -3, -2, -1, 0, 1, 2, 3, 4$




Рис. 3: Maps of F and G in the same notations as in figure 2.




Рис. 4: $f = \sqrt{1(z)} = \text{Factorial}^{-1/2}(z)$ by (10), left, and $f = \sqrt{\exp(z)}$, right, in the complex z -plane in the same notations as in figures 2, 3